



V1 serves as a motor cortex for visually guided saccades: evidence from human and monkey behavioral and neural data





Poster presentation at MütZü 2020 oculomotor research meeting, Tübingen, Germany Feb 14-15. 2020


Li Zhaoping, University of Tübingen, Max Planck Institute for Biological Cybernetics, Germany. www.lizhaoping.org li.zhaoping@tuebingen.mpg.de

Abstract: I proposed in the late 1990s and early 2000s that the primary visual cortex, V1, creates a bottom-up saliency map from visual inputs (Li 1999, 2002). This is the V1 Saliency Hypothesis (V1SH). According to V1SH, the highest V1 response to any visual location represents this location's saliency, defined as the strength of exogenous attentional attraction. Hence, the visual location evoking the largest V1 response is most likely to attract visual attention or evoke a gaze shift driven by bottom-up visual inputs. For example, a vertical bar in a background of horizontal bars is salient; so is a red dot among green dots, because the feature singletons evoke higher V1 responses than the homogeneous background items. Contextual influences via the intracortical horizontal connections make V1 neurons' responses dependent on contextual inputs outside the classical receptive field (e.g., Knierim & van Essen 1992). These contextual influences are such that neurons tuned to the same or similar features suppress each other, so that the neurons preferring and responding to the homogenous background inputs suppress each other's responses. The neurons responding to the feature singletons escape such suppression and the singletons are thus salient. Here I review some of the supporting evidence in human and monkey data from past works by my collaborators and myself. In lower vertebrates, the saliency map resides in the superior colliculus/optic tectum (Zhaoping 2016). More info. at <http://www.lizhaoping.org/zhaoping/V1Saliency.html>

V1 Saliency Hypothesis (V1SH, Li 1999, 2002): V1 creates a bottom-up saliency map to guide attentional/gaze shifts



V1SH prediction: Attention capture by an eye-of-origin singleton --- because V1 neurons are tuned to eye-of-origin, and V1 has iso-eye-of-origin suppression --- confirmed experimentally (Zhaoping 2008, 2012)



An ocular singleton, though irrelevant in a task to search for a uniquely oriented bar, pops out to attract attention and gaze shift, even though observers do not find it visually distinctive and are often unaware of its presence. This attraction prolongs reaction time (RT) to find the target. This observation (Zhaoping 2008, 2012), predicted by V1SH, is a fingerprint of V1, which is the only cortical area with a substantial number of neurons tuned to eye-of-origin of visual inputs.


In the dichoptic input above, the uniquely oriented bar is salient, because the V1 neuron tuned to and responding to it escapes iso-orientation suppression. The ocular (eye-of-origin) singleton is also salient, because the responding neuron escapes the iso-eye-of-origin suppression. In this example, the ocular singleton is task irrelevant but is more salient than the orientation singleton, attracting the first saccade in the search task.

References:


- Knierim JJ, van Essen DC. Neuronal responses to static texture patterns in area V1 of the alert macaque monkey. *J Neurophysiol.* 1992 Apr;67(4):961-80.
- Li Z. (1999) Contextual influences in V1 as a basis for pop out and asymmetry in visual search. *PNAS, USA* 96:10530-10535
- Li Z. (2002) A saliency map in primary visual cortex. *Trends in Cognitive Sciences.* 6:9-16
- Tehovnik, EJ, Slocum WM, Schiller PH (2003) Saccadic eye movements evoked by microstimulation of striate cortex. *Eur. J. of Neuroscience.* 17:870-878.
- Koene AR and Zhaoping L. (2007) Feature-specific interactions in salience from combined feature contrasts: Evidence for a bottom-up saliency map in V1. *Journal of Vision.* 7(7):6, 1-14,

V1SH prediction: given the same visual input, higher V1 responses (due to response fluctuations or neural stimulation) should lead to faster saccades to the feature singleton --- confirmed experimentally in monkeys (Yan et al. 2018, c.f. Tehovnik et al. 2003)

Monkey's task: start a trial by fixating at the central fixation point; after the bars appear, saccade to the orientation singleton (whose position is unpredictable) as quickly as possible



V1 neural responses to the orientation singleton (spikes/sec) before the saccade onset (usually > 200 ms)




V1 neurons' responses to the search target were measured while the monkey was doing the task. Usually it took more than 200 ms after stimulus onset before the saccadic onset. At around 40-50 ms after the stimulus onset, given the same visual inputs, trials with higher initial V1 neural responses to the target were more likely to evoke faster and accurate subsequent saccades to the target. This latency, at 40-60 ms, is so short that higher cortical areas or superior colliculus could not have provided this saliency signal to V1. This latency agrees with the time course of the contextual influences observed in Knierim and van Essen 1992. White et al. 2017 did not see this saliency signal at this short latency, but they saw a signal at a latency longer than that in superior colliculus, in their monkeys doing a similar task. Yan et al. 2018 showed that a signal at such a longer latency arises from top-down factors.


Tehovnik et al. 2003: microstimulation of V1 leads to saccades in monkeys to the receptive field's location

fMRI/ERP evidence: salient but imperceptible inputs (the cue below) activate early visual cortex but not the parietal cortex (Zhang, Zhaoping, Zhou, Fang, 2012)

An imperceptible cue that attracts attention




ERP and BOLD responses to this cue



V1SH fully predicts reaction times (RTs) using zero parameters, suggesting that V2 and other visual cortical areas do not contributing to saliency computation

(Koene & Zhaoping 2007, Zhaoping & Zhe 2015)



- Zhaoping L. (2008) Attention capture by eye of origin singletons even without awareness --- a hallmark of a bottom-up saliency map in the primary visual cortex. *Journal of Vision.* 8(5):1, 1-18,
- Zhang X., Zhaoping L., Zhou T. and Fang F. (2012) Neural activities in V1 create a bottom-up saliency map. *Neuron.* 73,183-192
- Zhaoping L. (2012) Gaze capture by eye of origin singletons: interdependence with awareness. *Journal of Vision.* 12(2):17, 1-22
- Zhaoping L. (2016) From the optic tectum to the primary visual cortex: migration through evolution of the saliency map for exogenous attentional guidance. *Current Opinion in Neurobiology.* 40:94-102.
- White BJ, Kan JY, Levy R, Itti L, Munoz DP. (2017) Superior colliculus encodes visual saliency before the primary visual cortex. *Proc Natl Acad Sci U S A.* 114(35):9451-9456.
- Yan Y., Zhaoping L., and Li W. (2018) Bottom-up saliency and top-down learning in the primary visual cortex of monkeys. *PNAS.* 115 (41) 10499-10504.